

## ACT Influenza Surveillance Report - No. 19, 2019

Summary: Influenza notifications have continued to decrease in week 38. If notifications continue to decline in the coming weeks, the 2019 influenza season will soon be over.

## Influenza notifications


Reporting Period: 1 January to 22 September 2019, Weeks 1-38

Influenza notifications have continued to decrease in the last week (Figure). Between 1 January and 22 September 2019, there were 3,813 notifications of influenza reported to ACT Health, with only 76 (2%) of these notifications occurring in the last week (Figure). Notifications in 2019 have surpassed the total number for 2017, noting that the 2019 influenza season began much earlier compared to previous seasons. In 2017, there were a total of 3,098 influenza notifications reported, and activity peaked at 327 notifications in one week (reporting week 33).

The proportion of influenza B notifications has increased in the last week compared to the previous weeks, with 29% of notifications in week 38 recorded as influenza B (compared with 23% in week 37). Year to date, 76% (n=2,893) of notifications were influenza A and 23% (n=876) were influenza B. There have been 44 notifications (1%) of cases co-infected with influenza A and B. Of the 359 (12%) influenza A notifications with subtype information available, 115 (32%) were A/H1 and 244 (68%) were A/H3.

In 2019, notification numbers overall have been highest among adults aged 20-64 years (49%, n=1,870), with only 12% (n=457) of notifications among children aged 0-4 years and 15% (n=562) among adults 65 years and over.

Figure. Number of influenza notifications, by week and year of onset, 1 January 2014 to 22 September 2019\*, ACT.



<sup>\*</sup>Data provided for the current and most recent weeks may be incomplete. All data are preliminary and subject to change as updates are received. Notification data include all cases of influenza diagnosed by a laboratory among residents of the ACT. Generally, notified cases represent only a small proportion of cases of influenza occurring in the community.